

TBE in Norway

Åshild K. Andreassen, Andrea P. Cotes-Perdomo, Arnulf Soleng,
Urusha Maharjan, Heidi H. Lindstedt, Kristian Alfsnes and Rikard Rykkvin

E-CDC risk status: endemic (last edited in June 2026, update for 2025: 97 reported cases)

History and current situation

In Norway, tick-borne encephalitis (TBE) has been a mandatory notifiable disease since 1975 (Norwegian Surveillance system for communicable diseases, MSIS).¹ According to ECDCs classification, coastal areas in southern Norway (counties of Agder, Vestfold and Telemark) are endemic for TBE. Further, the counties of Østfold, Akershus and Buskerud, and western and northern Norway to Brønnøy municipality are imperiled.²⁻¹²

The first reported case of TBE occurred in 1997 at Tromøy in Agder County.¹³ This is a region with holiday cabins and outdoor recreation areas for both local inhabitants and tourists, and it is known for high temperatures during spring and summer. In addition, TBE antibodies in dogs and tick-borne encephalitis virus (TBEV) in ticks have been detected in this area.^{10,12-15}

A total number of 519 TBE cases have been reported to MSIS as of February 2024 (Fig. 1). Of these, 420 cases are autochthonous infections, while 99 cases were infected abroad or have an unknown infection history. The number of cases varies annually between 1 and 86 (Table 1 and Fig. 1). Data for 2018 to 2023 shows an increase in the number of cases, especially in the counties of Vestfold and Telemark (Fig. 5, MSIS, February 2023). In 2023 the first case from Vestland County was reported to MSIS. The TBE patients' age distribution is in accordance with other European studies, with a higher infection rate for those older than 30 years (Table 2 and Fig. 2).¹⁶⁻¹⁷ According to MSIS, the reported cases in Norway are represented by the counties of Agder, Vestfold and Telemark, and Viken, all located in the southern part of the country (Fig. 3). No cases are reported from the northern coastal areas but a few cases are reported from the western areas and the area east of the Oslofjord, even though outdoor recreation activities are common in the whole country.

Ticks and TBEV in Norway

The castor bean tick (*Ixodes ricinus*) is the most common tick species in Europe,¹⁸ and considered to be the major vector of the European TBE-virus.¹⁹⁻²⁰ The geographical distribution of *I. ricinus* in Norway has been investigated in several studies.^{2-4,10,21-25} Both Tambs-Lyche (1943) and Mehl (1983) found *I. ricinus* to be mainly distributed in the coastal

areas of Norway, from the southeastern border to Sweden, along the southern and western coastline, up to Nordland County at ~65.1°N.^{2,3,21-25} The density of ticks varies between locations, even when separated by short distances. This is probably caused by differences in microclimatic conditions, vegetation, and density of vertebrate hosts. However, locations with a high density of ticks are found all over the major distributional range. The density of ticks declines rapidly with both increasing distance from the coast and increasing altitude. In a multi-source study, Jore et al. (2011) suggested that tick populations in Norway had undergone recent shifts in latitudinal and altitudinal range.²⁴ This result is, however, disputed in recent studies.^{4,5}

Although ticks are reported far outside (i.e. northeast) of the hitherto established distribution limit of *I. ricinus* in Norway, the vast majority of these are engorged females.²⁴⁻²⁶

Migratory birds may deposit engorged larvae or nymphs in areas where temperatures permit development to the next stage but not completion of the life cycle. Thus, such records do not constitute evidence for established and sustainable tick populations as this requires the presence of all the active stages (larvae, nymphs, and adults) in a locality for at least two consecutive seasons.²⁷⁻²⁸ Using flagging and dragging, Soleng et al. (2018) found tick larvae, nymphs and adults to be abundant at 64.5 and 65.1°N. Only a few tick nymphs and adults, and no larvae, were found at locations close to 66°N. At several locations from 66.3°N up to 67.5°N no ticks were found.³ In a recent study by Hvidsten et al. (2020), the occurrence of ticks in northern Norway was examined by dragging in 109 separate locations between the latitudes of 64°N and 70°N. The northernmost location with a permanent *I. ricinus* population was at 66.2°N on the Island of Dønna (Fig. 4).^{5,6} It is noteworthy that the taiga tick (*Ixodes persulcatus*) and the meadow tick (*Dermacentor reticulatus*) were not detected in a large screening of ticks collected in the southern part of Norway in 2016.²⁹

Studies of *I. ricinus* in Norway have detected TBEV in nymphs with prevalence ranging from 0% to 1.1%. In adult ticks collected from the same areas, the prevalence ranges from 0% to 20.6%. TBEV positive ticks have been found in sampling areas along the Norwegian coastline from the east of Østfold County to Brønnøy in Nordland County.¹² The

highest estimated TBEV prevalence in adult ticks has been found in the counties of Rogaland and Vestfold and Telemark. In nymphs, the highest prevalence has been found in Vestfold, Telemark, Agder and Rogaland.¹²

Historically, the first suggested TBEV isolate from Norway was collected in *I. ricinus* from Vestland County (former Sogn and Fjordane) in June 1976 as described by Traavik and co-workers. Five virus strains with a close serological relationship to the TBEV complex were detected in this study.²⁵

One pool of ten nymphs collected from southern Norway has been whole-genome sequenced and phylogenetically characterized. The strain, “Mandal 2009”, was found to belong to the Scandinavian group of the European TBEV subtype. Interestingly, “Mandal 2009” revealed a shorter form of the TBEV genome within the 3′ non-coding region, like the highly virulent “Hypr” strain.²⁰ Recent unpublished findings indicate circulation of at least one new TBEV variant in Norway from two new areas. This variant in the TBEV sequence is detected in a tick and one patient sample, both different from the previous Mandal 2009 strain.^{31,32}

Seroprevalence in animals

In addition to tick studies, a seroprevalence study has detected TBE antibodies in specimens from cervids (deer) collected in Farsund (Agder County) and Molde (Møre and Romsdal County). In Farsund, located on the southern coast of Norway, 41% (22 of 54 animals) were TBE-positive. This contrasts with Molde, situated midwest, where the prevalence was 1.6% (1 of 64 animals). The same study detected antibodies to Louping ill virus (LIV), a closely related flavivirus, in 14.8% (8 of 54) of the analyzed cervid sera from Farsund.²¹

A recent seroprevalence study of cervids where serum samples were collected across Norway found TBEV antibodies in the municipalities of Steinkjer, Vindafjord, Søgne, Birkenes, Lardal, Larvik and Halden (Fig. 4). The overall seroprevalence was 4.6%. Antibodies against TBEV detected by serum neutralization test were present in 9.4% of the moose samples, 1.4% in red deer, 0.7% in roe deer, and 0% in reindeer.¹¹

Ticks (6850 nymphs and 765 adults) from eastern, western, and northern Norway were analyzed for LIV using an in-house real-time polymerase chain reaction (PCR), none of these were positive (unpublished data). However, a recent study by Ytrehus et al. detected antibodies against LIV in willow ptarmigan (*Lagopus lagopus lagopus*) across the whole country. The study suggested that either LIV or a cross-reacting virus infects ptarmigan in Norway, also at high altitudes and latitudes.²⁷

There is limited knowledge of TBEV in domestic animals in Norway. A recent study reported TBEV RNA in

unpasteurized cow milk from three farms located in southern and northern Norway in 5.4% of the tested animals. Seropositive animals were only detected at one farm in southern Norway, in 88.2% of the tested animals.¹⁰ This is higher than in a previous study by Traavik (1973), where a seroprevalence of 17.7% was detected in bovine sera in western Norway.²²

Seroprevalence in humans

In Søgne municipality, a TBE endemic area of southern Norway, a TBEV seroprevalence of 3.1% (45/1,453) was found in the general adult population. Among individuals not vaccinated against TBEV and/or yellow fever, the seroprevalence of IgG antibodies to TBEV was 1.4% (6/419).³³ A recent blood donor study from TBE endemic areas in Vestfold and Telemark found a low seroprevalence of 0.4% (4/1,123). Out of the 1,123 analyzed samples, 21 had neutralizing antibodies to TBEV, of which 17 reported a previous TBE vaccination.³²

Three seroprevalence studies in humans from presumed non-endemic areas have been published. Larsen et al. detected TBE immunoglobulin G (IgG) antibodies among 0.65% of blood donors in Viken County (former Østfold) in southeastern Norway.²³ The second study in 1,213 blood donors was performed in Vestland County (former Sogn and Fjordane), located in western Norway. TBE IgG antibodies (ELISA) were detected in five (0.4%) of these samples. However, four of these were reported to be vaccinated against flaviviruses and one was negative by neutralization test.²⁴ In 1979, Traavik detected a 19.6% seroprevalence from Vestland County. However, these results were not confirmed with a neutralization test and thus, may be explained by cross-reactions to LIV, vaccine-related flaviviruses, or nonspecific binding in the test.²⁶

TBEV in ticks in Norway is widely distributed (Fig. 4). It has been a puzzle why there have been no reports of patients outside the endemic areas. However, this seems to undergo a change with increasing incidence and the geographical expansion of cases towards north and east as illustrated (Appendix Fig. 1; Fig. 5).

Conclusion

In summary, TBE is endemic in parts of Norway and the number of human TBE cases has been increasing in recent years. Clinical TBE cases are only found in southern parts of Norway; however, the results from both prevalence studies in ticks and seroprevalence studies in humans and animals indicate that TBEV might be widespread in the country, and not limited to the southern region. This is highly relevant information for public health considerations and risk evaluation. Further studies on tick distribution and prevalence of TBEV in ticks, humans and animals in Norway are currently ongoing.

Overview of TBE in Norway

Table 1: Virus, vector, transmission of TBE in Norway

Viral subtypes, distribution ^{3,7,8,10-12,21,25}	Western subtype. TBEV is distributed in <i>Ixodes ricinus</i> ticks in the following counties: Viken, Vestfold and Telemark, Agder, Rogaland, Vestland, Møre og Romsdal, Trøndelag and Nordland. Human TBE cases have been reported in the following counties: Agder, Vestfold and Telemark, and Viken. Source: www.fhi.no Norwegian Surveillance System for Communicable Diseases (MSIS)
Reservoir animals	Small rodents in the genera <i>Apodemus</i> and <i>Myodes</i> . ²⁶
Infected tick species (%)	<i>Ixodes ricinus</i> (0–1.1% in nymphs and 0–20.6% in adults). ¹⁰
Dairy product transmission	Not documented. TBEV RNA has been detected in unpasteurized cow milk.⁸

Table 2: TBE-reporting and vaccine prevention in Norway


Mandatory TBE-reporting	Hospitals and General Practitioners Only cases with meningitis/encephalitis are notifiable. Criteria: - Detection of specific antibody response in serum and/or cerebrospinal fluid and/or - Detection of TBEV in cerebrospinal fluid and/or serum by isolation and nucleic acid analysis Source: www.fhi.no
Other TBE-Surveillance	Ongoing studies: The Barents region project (ID B1710). Emerging infections. Capacity building on vector-borne infections in the Barents Region, Norway and Russia. EMERGING VIRUS: Vector-borne infections in Norway – Understanding the emergence of viral vector-borne diseases in a One Health perspective by studies of dynamics, distribution, climate, genetic diversity, biogeographic distribution, risk of infection, surveillance and diagnosis Observation and prognosis of TBE-patients (Telemark Sykehus HF). North-Tick (Interreg VB, North Sea program): Vector-borne pathogens/infections in the North Sea area. Experimental infection study of sheep with TBEV (ref): Transmission to lambs via milk (ref) TBFV net (EEA-project): surveillance and research on tick-borne flaviviruses Development of pipeline of whole genome sequencing of TBEV (ref)
Special clinical features	No. TBE has been mandatorily notifiable to MSIS (Norwegian Surveillance System for Communicable Diseases) since 1975. Source: www.fhi.no
Available vaccines	TicoVac, Pfizer TicoVac Junior, Pfizer Source: <i>The Norwegian Medicines Agency</i>
Vaccination recommendations and reimbursement	People who spend a lot of time outdoors in areas with a known risk of TBE infection (Table 1) should consider TBE vaccination. This recommendation applies especially to people who often get tick bites such as forestry workers, orienteers, and others who spend much time outdoors. There is no reimbursement in place. Source: www.fhi.no

Table 2: TBE-reporting and vaccine prevention in Norway

Vaccine uptake by age group/risk group/general population	In Norway, all immunizations should be registered into the national immunization register, SYSVAK. According to SYSVAK for 2024, 103 749 persons have been vaccinated against TBE and there were sold 172 355 doses which is an increase from 2022 and 2023. There is no information about risk factors in the register. For vaccines outside the childhood immunization program, registration into SYSVAK is consensual. The number of doses actually given therefore could be higher than the numbers registered, and this is probably the case for TBE vaccine, as the number of doses distributed is higher than the number of doses entered into the register. <i>Source: Norwegian Immunization Registry (SYSVAK)</i>
Name, address/ website of TBE NRC	Norwegian Institute of Public Health. <i>Source: www.fhi.no</i>


Figure 1: Burden of TBE in Norway 1994-2024*

(last edited in May 2026, update for 2025: 97 reported cases)




*data per February 2025 (MSIS).

Source Data: Appendix—Figure 1

Figure 2: Age and gender distribution of TBE in Norway 1994–2025*

*data per February 2025 (MSIS).

Source Data: Appendix—Figure 2

Figure 3: TBE cases in Norway 1994–2024 (MSIS)

Datakilde skogflåttencefalitt (TBE): MSIS 1994-2024
 Kartet er laget i R versjon 4.4.0, osmaps 2024.5.27 (data fra Kartverket via Geonorge)
 Folkehelseinstituttet, november, 2025

Source: <https://www.fhi.no/nettpub/smittevernveilederen/sykdommer-a-a/skogflattencefalitt-tbe-virusinfeks/#forekomst-i-norge>

Figure 4: Geographical locations where tick-borne encephalitis virus has been detected in *Ixodes ricinus* in Norway from 2004 to 2020:
 ○ No ticks found, ● Ticks with TBEV, ● TBEV antibodies in animals, ● TBEV in ticks, cow milk, and TBEV antibodies in animals
 ● Site where *Ixodes persulcatus* have been detected 2023.^{3, 7-10, 19, 21, 25}



Map from © Kartverket (<https://www.kartverket.no/> Attribution 4.0 International (CC BY 4.0))

Figure 5: Expanded geographical distribution of reported TBE cases in Norway between 2009 and 2025. The red line shows the distribution border in 2009, the areas north and east of this, represents areas where new cases have been reported after 2009.




Table 3: Overview of monthly reported TBE cases 2025

Month	Infected in Norway	Unknown infection site	Infected abroad	Total 2025	Total 1994-2025
January		1		1	2
April	3			3	4
May	5	1	1	10	42
June	9	3		12	75
July	14	4	1	19	128
August	17		2	19	167
September	12			12	139
October	12	1		13	99
November	7			7	31
December	1			1	5
Total 2025	83	10	4	97	692

Appendix

Source data: Figure 1

Year	Number of cases	Incidence / 10 ⁵
1994	2	<0.1
1995	0	0
1996	0	0
1997	0	0
1998	1	<0.1
1999	1	<0.1
2000	1	<0.1
2001	0	0
2002	2	<0.1
2003	1	<0.1
2004	4	<0.1
2005	4	<0.1
2006	5	0.1
2007	13	0.2
2008	11	0.2
2009	10	0.2
2010	11	0.2
2011	14	0.3
2012	7	0.1
2013	6	0.1
2014	13	0.2
2015	9	0.2
2016	12	0.2
2017	16	0.3

Year	Number of cases	Incidence / 10 ⁵
2018	26	0.5
2019	35	0.7
2020	41	0.8
2021	71	1.3
2022	90	1.6
2023	113	2.0
2024	76	1.4
2025	97	1.7

Source data: Figure 2

Age group (years)	Females	Males	All
0-9	8	13	21
10-19	23	31	54
20-29	18	21	39
30-39	29	53	82
40-49	44	83	127
50-59	60	68	128
60-69	37	83	120
>70	45	76	121

Acknowledgements

We thank Trude Marie Lyngstad for making Figure 3.

Authors and affiliations

Åshild K. Andreassen, Andrea P. Cotes-Perdomo, Arnulf Soleng, Urusha Maharjan, Heidi H. Lindstedt and Rikard Rykkvin

Norwegian Institute of Public Health, Division of Infection Control and Environmental Health

Kristian Alfsnes

Norwegian Institute of Public Health, Department of Bacteriology

Contact: ashildkristine.andreassen@fhi.no

Citation:

Andreassen KA, Cotes-Perdomo AP, Soleng A, Maharjan U, Lindstedt HH, Alfsnes K, Rykkvin R. TBE in Norway. Chapter 13. In: Dobler G, Erber W, Bröker M, Chitimia-Dobler L, Schmitt HJ, eds. *The TBE Book*. 9th ed. Singapore: Global Health Press; 2026. doi:10.33442/26613980_13-24-9

References

1. Norwegian Surveillance System for Communicable Diseases (MSIS). 2021.
2. Tambs-Lyche. *Ixodes ricinus* og piroplasmosen i Norge (Meddelelse fra Bergen Museums zoologiske avdeling). 1943;55:337-366.
3. Mehl R. The distribution and host relations of Norwegian ticks (Acari, Ixodidae). *Fauna norvegica series B*. 1983;30:4651.
4. Soleng A, et al. Distribution of *Ixodes ricinus* ticks and prevalence of tick-borne encephalitis virus among questing ticks in the Arctic Circle region of northern Norway. *Ticks Tick Borne Dis*. 2018;9(1):97-103. doi:10.1016/j.ttbdis.2017.10.002
5. Hvidsten D, Frafjord K, Gray JS, Henningsson A, Jenkins A, Kristiansen BE, Lager M, Rognerud B, Slåtsve AM, Stordal F, Stuen S, Wilhelmsson P. The distribution limit of the common tick, *Ixodes ricinus*, and some associated pathogens in northwestern Europe. *Ticks Tick Borne Dis*. 2020;3:101388. doi:10.1016/j.ttbdis.2020.101388
6. Hvidsten D, Stuen S, Jenkins A, Dienus O, Olsen RS, Kristiansen BE, Mehl R, Matussek A. *Ixodes ricinus* and *Borrelia* prevalence at the Arctic Circle in Norway. *Ticks Tick Borne Dis*. 2014;5(2):107-12. doi:10.1016/j.ttbdis.2013.09.003
7. Jore S, et al. Multi-source analysis reveals latitudinal and altitudinal shifts in range of *Ixodes ricinus* at its northern distribution limit. *Parasit Vectors*. 2011;4:84.
8. Jenkins A, Hvidsten D, Matussek A, Lindgren PE, Stuen S, Kristiansen BE. *Borrelia burgdorferi sensu lato* in *Ixodes ricinus* ticks from Norway: evaluation of a PCR test targeting the chromosomal *flaB* gene. *Exp Appl Acarol*. 2012; 58(4):431-9. doi:10.1007/s10493-012-9585-2
9. Paulsen KM, Pedersen BN, Soleng A, Okbaldet YB, Pettersson JH, Dudman SG, Ottesen P, Vik IS, Vainio K, Andreassen A. Prevalence of tick-borne encephalitis virus in *Ixodes ricinus* ticks from three islands in north-western Norway. *APMIS*. 2015; 123(9):759-64. doi: 10.1111/apm.12412
10. Paulsen KM, Stuen S, das Neves CG, Suhel F, Gurung D, Soleng A, Stiasny K, Vikse R, Andreassen AK, Granquist EG. Tickborne encephalitis virus in cows and unpasteurized cow milk from Norway. *Zoonoses Public Health*. 2019;66(2):216-222. doi: 10.1111/zph.12554
11. Paulsen KM, das Neves CG, Granquist EG, Madslien K, Stuen S, Pedersen BN, Vikse R, Rocchi M, Laming E, Stiasny K, Andreassen AK. Cervids as sentinel-species for tick-borne encephalitis virus in Norway - A serological study. *Zoonoses Public Health*. 2019. doi:10.1111/zph.12675.
12. Vikse R, Paulsen KM, Edgar KS, Pettersson JH-O, Ottesen PS, Okbaldet YB, Kiran N, Lamsal A, Lindstedt HEH, Pedersen BN, Soleng A, Andreassen AK. Geographical distribution and prevalence of tick-borne encephalitis virus in questing *Ixodes ricinus* ticks and phylogeographic structure of the *Ixodes ricinus* vector in Norway. *Zoonoses Public Health*. 2020:1-12. doi:10.1111/zph.12696.
13. Skarpaas T, Ljøstad U, Sundøy A. First human cases of tickborne encephalitis, Norway. *Emerg Infect Dis*. 2004;10(12):2241-3. doi:10.3201/eid1012.040598.
14. Skarpaas T, Golovljova I, Vene S, Ljøstad U, Sjørnsen H, Plyusnin A, Lundkvist A. Tickborne encephalitis virus, Norway and Denmark. *Emerg Infect Dis*. 2006;12(7):1136-8. doi:10.3201/eid1207.051567
15. Csángó PA, Blakstad E, Kirtz GC, Pedersen JE, Czettel B. Tickborne encephalitis in southern Norway. *Emerg Infect Dis*. 2004;10(3):533-4. doi:10.3201/eid1003.020734
16. Andreassen A, Jore S, Cuber P, Dudman S, Tengs T, Isaksen K, Hygen HO, Viljugrein H, Anestad G, Ottesen P, Vainio K. Prevalence of tick borne encephalitis virus in tick nymphs in relation to climatic factors on the southern coast of Norway. *Parasit Vectors*. 2012;5:177. doi:10.1186/1756-3305-5-177.
17. Norwegian Surveillance System for Communicable Diseases (MSIS). 2019; Available from: www.MSIS.no, 31.10.19.
18. Radzisauskiene D, Zagminas K, Asokliene L, Jasionis A, Mameniskiene R, Ambrozaitis A, Jancoriene L, Jatuzis D, IPetraityte I, Mockiene E. Epidemiological patterns of tickborne encephalitis in Lithuania and clinical features in adults in the light of the high incidence in recent years: a retrospective study. *Eur J Neurol*. 2018;25(2):268-74. collected from recreational islands in southern Norway. *Ticks Tick Borne Dis*. 2018;9(5):1098-1102. doi:10.1111/ene.13486
19. Kaminski M, Grummel V, Hoffmann D, Berthele A, Hemmer B. The spectrum of aseptic central nervous system infections in southern Germany - demographic, clinical and laboratory findings. *Eur J Neurol*. 2017;24(8):1062-70. doi:10.1111/ene.13335
20. Asghar N, Lindblom P, Melik W, Lindqvist R, Haglund M, Forsberg P, Overby AK, Andreassen A, Lindgren PE, Johansson M. Tick-borne encephalitis virus sequenced directly from questing and blood-feeding ticks reveals quasispecies variance. *PLoS One*. 2014;9(7):e103264. doi:10.1371/journal.pone.0103264
21. Ytrehus B, Vainio K, Dudman SG, Gilray J, Willoughby K. Tickborne encephalitis virus and louping-ill virus may cocirculate in Southern Norway. *Vector Borne Zoonotic Dis*. 2013;13(10):762-8. doi:10.1089/vbz.2012.1023
22. Traavik T. Serological investigations indicating the existence of tick-borne encephalitis virus foci along the Norwegian coast. *Acta Pathol Microbiol Scand B Microbiol Immunol*. 1973;81(1):138-42. doi:10.1111/j.1699-0463.1973.tb02197.x
23. Larsen AL, Kanestrom A, Bjorland M, Andreassen A, Soleng A, Vene S, Dudman SG. Detection of specific IgG antibodies in blood donors and tick-borne encephalitis virus in ticks within a non-endemic area in southeast Norway. *Scand J Infect Dis*. 2014;46(3):181-4. doi: 10.3109/00365548.2013.865140.
24. Hjetland R, Henningsson AJ, Vainio K, Dudman SG, Grude N, Ulvestad E. Seroprevalence of antibodies to tick-borne encephalitis virus and *Anaplasma phagocytophilum* in healthy adults from western Norway. *Infect Dis (Lond)*. 2015;47(1):52-6. doi:10.3109/00365548.2014.959044

25. Traavik T, Mehl R, Wiger R. The first tick-borne encephalitis virus isolates from Norway. *Acta Pathol Microbiol Scand B*. 1978;86(4):253-5. doi:10.1111/j.1699-0463.1978.tb00040.x
26. Traavik T. Antibodies to tick-borne encephalitis virus in human sera from the western coast of Norway. *Acta Pathol Microbiol Scand B*. 1979;87B(1):9-13. doi:10.1111/j.1699-0463.1979.tb02396.x
27. Kjelland V, Paulsen KM, Rollum R, Jenkins A, Stuen S, Soleng A, Edgar KS, Lindstedt HH, Vaino K, Gibory M, Andreassen AK. Tick-borne encephalitis virus, *Borrelia burgdorferi sensu lato*, *Borrelia miyamotoi*, *Anaplasma phagocytophilum* and *Candidatus Neoehrlichia mikurensis* in *Ixodes ricinus* ticks collected from recreational islands in southern Norway. *Ticks Tick Borne Dis*. 2018;9(5):1098-1102.
28. Jaenson TG, Hjertqvist M, Bergstrom T, Lundkvist A. Why is tick-borne encephalitis increasing? A review of the key factors causing the increasing incidence of human TBE in Sweden. *Parasit Vectors*. 2012;5:184. doi:10.1186/17563305-5-184
29. Cotes-Perdomo A, Soleng A, Alfsnes K, Andreassen Å. First report of the taiga tick *Ixodes persulcatus* in Norway. *Ticks and Tick-borne Diseases*. 2025;16(4):102508. doi:10.1016/j.ttbdis.2025.102508
30. Maharjan U, Skudal HK, Asghar N, et al. Novel variants of tickborne encephalitis virus from patient and tick samples in Norway. *Ticks and Tick-borne Diseases*. 2025;16(4):102501. doi:10.1016/j.ttbdis.2025.102501
31. Holten AR, Bjerkhaug AW, Maharjan U, et al. A case of fatal hemorrhagic fever associated with tick-borne encephalitis virus infection. *International Journal of Infectious Diseases*. 2025;159:107989. doi:10.1016/j.ijid.2025.107989
32. Thortveit ET, Aase A, Petersen LB, Lorentzen ÅR, Mygland Å, Ljøstad U. Human seroprevalence of antibodies to tickborne microbes in southern Norway. *Ticks Tick Borne Dis*. 2020;11(4):101410. doi: 10.1016/j.ttbdis.2020.101410.
33. Ytrehus B, Rocchi M, Brandsegg H, Turnbull D, Miller A, Pedersen HC, Kålås JA, Nilsen EB. Louping-ill virus serosurvey of willow ptarmigan (*Lagopus lagopus lagopus*) in Norway. *J Wildl Dis*. 2021 Apr 1;57(2):282-291. doi: 10.7589/JWD-D-2000068.